*Fluid Phase Equilibria*, 43 (1988) 363–364
Elsevier Science Publishers B.V., Amsterdam – Printed in The Netherlands



## Contents Volume 43 (1988)

*Volume 43, No. 1*

Solubility of solids in supercritical solvents. IV. Mean-field lattice gas description for the *p-T-x* space diagram of the system ethylene–naphthalene
R. van der Haegen, R. Koningsveld, L. van Opstal (Wilrijk, Belgium) and L.A. Kleintjens (Geleen, The Netherlands) . . . . . 1–19

Prediction of activity coefficients of electrolytes in aqueous mixed electrolyte solutions
A. Kumar (Pune, India) . . . . . 21–27

Vapor–liquid equilibria of the binary mixtures methyl chloride–methanol and dimethyl ether–methyl chloride and of the ternary mixture dimethyl ether–methyl chloride–methanol
H. Holldorff and H. Knapp (Berlin, F.R.G.) . . . . . 29–44

Solubility of monocrotaline in supercritical carbon dioxide and carbon dioxide–ethanol mixtures
S.T. Schaeffer, L.H. Zalkow and A.S. Teja (Atlanta, GA, U.S.A) . . . . . 45–56

Vapor–liquid equilibrium calculations of multicomponent systems based on Taylor series approximation for quasichemical equilibria
D. Brennecke and R. Lacmann (Braunschweig, F.R.G.) . . . . . 57–71

Solute–solvent interactions: dissolution of sparingly soluble silver salts of dibasic and tribasic inorganic and organic acids in water + acetic acid mixtures
U.N. Dash and E. Pattanaik (Bhubaneswar, India) . . . . . 73–76

Vapor–liquid equilibria of 2-propanol + water + *N,N*-dimethyl formamide
H.S. Wu, D. Hagewiesche and S.I. Sandler (Newark, DE, U.S.A.) . . . . . 77–89

Isobaric vapour–liquid equilibria of ethanol–toluene–sodium acetate mixtures at various system pressures
T.C. Tan and H.C. Ti (Kent Ridge, Republic of Singapore) . . . . . 91–104

Phase equilibria for four binary systems containing propylene
K. Ohgaki, H. Takata, T. Washida and T. Katayama (Osaka, Japan) . . . . . 105–113

*Guide for Authors* . . . . . 115–118

*Announcement* . . . . . 119

*Volume 43, Nos. 2–3*

Correlation of equation of state parameters for the associated perturbed anisotropic chain theory
P. Vimalchand, G.D. Ikonomou and M.D. Donohue (Baltimore, MD, U.S.A.) 121–135

Prediction of vapor–liquid equilibria for electrolyte solution by the extended Pitzer (1980) model. Part 1. Physical interpretation of the model parameter
X.H. Lu, Y.R. Wang and J. Shi (Nanjing, People's Republic of China) . . . . . 137–152

Estimation of the second cross virial coefficient from complete vapour–liquid equilibrium data. A simulation study
V. Dohnal (Prague, Czechoslovakia) . . . . . 153–170

A new approach for calculating the accessible volume in equations of state for mixtures. I. Theory and implementation in the van der Waals equation of state
B. Smit and K.R. Cox (Amsterdam, The Netherlands) . . . . . 171–180
A new approach for calculating the accessible volume in equations of state for mixtures. II. Application to Lennard–Jones mixtures
B. Smit and K.R. Cox (Amsterdam, The Netherlands) . . . . . 181–189
Methods for predicting and correlating the vapour pressures of pure compounds
A. Vetere (Milan, Italy) . . . . . 191–203
Statistical mechanical test of MHEMHS model of the interaction third virial coefficients
E.Z. Hamad, G.A. Mansoori (Chicago, IL, U.S.A.) and K.R. Hall (College Station, TX, U.S.A.) . . . . . 205–212
Description of the phase equilibrium of sulfuric acid with the NRTL equation and a solvation model in a wide concentration and temperature range
A. Bosen and H. Engels (Aachen, F.R.G.) . . . . . 213–230
Group contribution methods for molecular mixtures. I. Interaction site models
L.L. Lee (Norman, OK, U.S.A.) and J.M. Haile (Clemson, SC, U.S.A.) . . . . . 231–261
$P-V-T$ data and virial coefficients for gaseous methane–water mixtures with correction for adsorption effects
L.L. Joffrion and P.T. Eubank (College Station, TX, U.S.A.) . . . . . 263–294
Analysis of excess enthalpies of ester + 1-chloroalkanes with two group contribution models: primary parameters
J. Ortega, J.S. Matos (Canarias, Spain) M.I. Paz-Andrade, J. Fernandez and L. Pias (Santiago, Spain) . . . . . 295–316
Quaternary liquid–liquid equilibrium: water–ethanol–2-ethylhexanol–decalin at 25 °C. Unusual shape of the solubility surface
F. Ruiz, V. Gomis and R.F. Botella (Alicante, Spain) . . . . . 317–327
Extraction of 1-methylnaphthalene and *m*-cresol with supercritical carbon dioxide and ethane
R.J. Lee and K.C. Chao (West Lafayette, IN, U.S.A.) . . . . . 329–340
Thermodynamics of some ternary mixtures of non-electrolytes at 308.15 K
H.P. Dahiya, P.P. Singh and S. Dagar (Rohtak, India) . . . . . 341–354

*Short Communication*
Limitations of differential static apparatus in dilute regions
P. Alessi, M. Fermeglia and I. Kikic (Trieste, Italy) . . . . . 355–358

*Errata* . . . . . 359–360

*Author Index* . . . . . 361

*Contents of Volume 43 (1988)* . . . . . 363–364